

"melt" in the order of their stability, lowest to highest. When a region of lower stability is near the end of a segment of duplex, and the next region of complementarity along the strand has a higher melting temperature, a temperature can be found that will cause the terminal region of duplex to melt first, opening the first bubble, and thereby creating a preferred substrate structure of the cleavage by the 5' nucleases of the present invention (Figure 40a). The site of such cleavage would be expected to be on the 5' arm, within 2 nucleotides of the junction between the single and double-stranded regions (Lyamichev *et al.*, *supra.* and U.S. Patent No. 5,422,253)

An additional oligonucleotide could be introduced to basepair along the target nucleic acid would have a similar effect of opening this bubble for subsequent cleavage of the unpaired 5' arm (Figure 40b and Figure 6). Note in this case, the 3' terminal nucleotides of the upstream oligonucleotide anneals along the target nucleic acid sequence in such a manner that the 3' end is located within the "bubble" region. Depending on the precise location of the 3' end of this oligonucleotide, the cleavage site may be along the newly unpaired 5' arm, or at the site expected for the thermally opened bubble structure as described above. In the former case the cleavage is not within a duplexed region, and is thus not invasive cleavage, while in the latter the oligonucleotide is merely an aide in inducing cleavage at a site that might otherwise be exposed through the use of temperature alone (*i.e.*, in the absence of the additional oligonucleotide), and is thus not considered to be invasive cleavage.

In summary, any arrangement of oligonucleotides used for the cleavage-based detection of a target sequence can be analyzed to determine if the arrangement is an invasive cleavage structure as contemplated herein. An invasive cleavage structure supports cleavage of the probe in a region that, in the absence of an upstream oligonucleotide, would be expected to be basepaired to the target nucleic acid.

Example 26 below provides further guidance for the design and execution of a experiments which allow the determination of whether a given arrangement of a pair of upstream and downstream (*i.e.*, the probe) oligonucleotides when annealed along a target nucleic acid would form an invasive cleavage structure.

## V. Fractionation Of Specific Nucleic Acids By Selective Charge Reversal

Some nucleic acid-based detection assays involve the elongation and/or shortening of oligonucleotide probes. For example, as described herein, the primer-directed, primer-independent, and invader-directed cleavage assays, as well as the "nibbling" assay all involve the cleavage (*i.e.*, shortening) of oligonucleotides as a means for detecting the presence of a target nucleic sequence. Examples of other detection assays which involve the shortening of an oligonucleotide probe include the "TaqMan" or nick-translation PCR assay described in U.S. Patent No. 5,210,015 to Gelfand *et al.* (the disclosure of which is herein incorporated by reference), the assays described in U.S. Patent Nos. 4,775,619 and 5,118,605 to Urdea (the disclosures of which are herein incorporated by reference), the catalytic hybridization amplification assay described in U.S. Patent No. 5,403,711 to Walder and Walder (the disclosure of which is herein incorporated by reference), and the cycling probe assay described in U.S. Patents Nos. 4,876,187 and 5,011,769 to Duck *et al.* (the disclosures of which are herein incorporated by reference). Examples of detection assays which involve the elongation of an oligonucleotide probe (or primer) include the polymerase chain reaction (PCR) described in U.S. Patent Nos. 4,683,195 and 4,683,202 to Mullis and Mullis *et al.* (the disclosures of which are herein incorporated by reference) and the ligase chain reaction (LCR) described in U.S. Patent Nos. 5,427,930 and 5,494,810 to Birkenmeyer *et al.* and Barany *et al.* (the disclosures of which are herein incorporated by reference). The above examples are intended to be illustrative of nucleic acid-based detection assays that involve the elongation and/or shortening of oligonucleotide probes and do not provide an exhaustive list.

Typically, nucleic acid-based detection assays that involve the elongation and/or shortening of oligonucleotide probes require post-reaction analysis to detect the products of the reaction. It is common that, the specific reaction product(s) must be separated from the other reaction components, including the input or unreacted oligonucleotide probe. One detection technique involves the electrophoretic separation

of the reacted and unreacted oligonucleotide probe. When the assay involves the cleavage or shortening of the probe, the unreacted product will be longer than the reacted or cleaved product. When the assay involves the elongation of the probe (or primer), the reaction products will be greater in length than the input. Gel-based electrophoresis of a sample containing nucleic acid molecules of different lengths separates these fragments primarily on the basis of size. This is due to the fact that in solutions having a neutral or alkaline pH, nucleic acids having widely different sizes (*i.e.*, molecular weights) possess very similar charge-to-mass ratios and do not separate [Andrews, Electrophoresis, 2nd Edition, Oxford University Press (1986), pp. 153-154]. The gel matrix acts as a molecular sieve and allows nucleic acids to be separated on the basis of size and shape (*e.g.*, linear, relaxed circular or covalently closed supercoiled circles).

Unmodified nucleic acids have a net negative charge due to the presence of negatively charged phosphate groups contained within the sugar-phosphate backbone of the nucleic acid. Typically, the sample is applied to gel near the negative pole and the nucleic acid fragments migrate into the gel toward the positive pole with the smallest fragments moving fastest through the gel.

The present invention provides a novel means for fractionating nucleic acid fragments on the basis of charge. This novel separation technique is related to the observation that positively charged adducts can affect the electrophoretic behavior of small oligonucleotides because the charge of the adduct is significant relative to charge of the whole complex. In addition, to the use of positively charged adducts (*e.g.*, Cy3 and Cy5 amidite fluorescent dyes, the positively charged heterodimeric DNA-binding dyes shown in Fig. 66, etc.), the oligonucleotide may contain amino acids (particularly useful amino acids are the charged amino acids: lysine, arginine, aspartate, glutamate), modified bases, such as amino-modified bases, and/or a phosphonate backbone (at all or a subset of the positions). In addition as discussed further below, a neutral dye or detection moiety (*e.g.*, biotin, streptavidin, etc.) may be employed in place of a positively charged adduct in conjunction with the use of amino-modified bases and/or a complete or partial phosphonate backbone.